THE LEADING
CAUSE OF
CANCER DEATH
IN CHILDREN
UNDER THE AGE
OF 5

MOST COMMON SOLID TUMOR IN CHILDREN UNDER THE

AGE OF 5

CHILDREN WHO
SURVIVED HAVE
A HIGH CHANCE
OF DEVELOPING
LONG TERM
SIDE EFFECTS
AS A RESULT OF
TREATMENT
THAT SAVED
THEIR LIVES.

ONLY 1
OUT OF 2
KIDS
DIAGNOSED
WILL
SURVIVE.

THE FACTS
ABOUT

NEUROBLASTOMA

SOME EARLY SYMPTOMS INCLUDE FEVER, NAUSEA, LOSS OF APPETITE AND PAIN, ALONG WITH BALANCE PROBLEMS & UNCONTROLLED MOVEMENTS.

> AN AGGRESSIVE CANCER THAT USUALLY STARTS IN THE ABDOMEN.

- * NEUROBLASTOMA ATTACKS THE NERVOUS SYSTEM & NEUROENDOCRINE GLANDS.
- * THE SYMPTOMS CAN VARY ACCORDING TO THE LOCATION OF THE TUMOR.

TREATMENT IS OFTEN **INTENSE, INVASIVE & PAINFUL** AND INCLUDES

• SURGERY • STEM CELL HARVESTS • RADIATION
• IMMUNOTHERAPY • MULTIPLE ROUNDS OF CHEMOTHERAPY